

# Sparrowhawk

*Accipiter nisus*

Category A

Breeding resident, winter visitor and passage migrant.

Knight and Tolputt included the Sparrowhawk in their “list of birds observed in Folkestone and its immediate neighbourhood” (1871). This list covered an area of six mile radius from Folkestone town hall. Whilst this range extends further inland than the current Folkestone and Hythe area, it is likely that it was present locally, for both Ticehurst (1909) and Harrison (1953) considered it to be a common breeding bird in most of the wooded districts of the county, though less so in the more sparsely wooded areas of the east and south-east.



Sparrowhawk at Saltwood Castle (Brian Harper)

Taylor *et al* (1981) however described a dramatic change in its fortunes and considered that “the Sparrowhawk has probably suffered more at the hands of man than any other species on the county list, and undoubtedly one of the more serious threats has been pesticide contamination during the 1950s and early 1960s, although keeping and woodland clearance will have also played a part. At this time numbers reached an all-time low and there was no confirmation of successful breeding between 1953 and 1974”.

Fortunately Henderson & Hodge (1998) were able to report that “the subsequent recovery in fortunes of this agile raptor has been remarkable. There was confirmation of six pairs nesting in Kent in 1975 and since then there has been a steady increase and eastward expansion, resulting in Sparrowhawks being present in ... 21% of all tetrads” in the 1988-94 survey. The expansion has since continued and Clements *et al* (2015) described it as now widespread in Kent, nesting in woodland, hedgerows and even in town centres.

## Breeding distribution

Figure 1 shows the breeding distribution by tetrad based on the results of the 2007-13 BTO/KOS atlas fieldwork.




Figure 1: Breeding distribution of Sparrowhawk at Folkestone and Hythe by tetrad (2007-13 BTO/KOS Atlas)

Breeding was confirmed in one tetrad, with probable breeding in four more and possible breeding in 16 others. The table below shows how this compares to previous atlases (Taylor *et al* 1981, Henderson & Hodge 1998). The confirmed and probable breeding categories have been combined to account for differing definitions of these in the first atlas.

Breeding atlases	1967 - 1973		1988 - 1994		2007 – 2013	
Possible	-	-	4	(13%)	16	(52%)
Probable/Confirmed	-	-	4	(13%)	5	(16%)
Total	-	-	8	(26%)	21	(68%)

The results of the first atlas were not published due to the species scarcity at the time, with it being only noted in 2% of the tetrads in the county (probable breeding in just two tetrads, with possible breeding in a further 17) so it was most likely to have been absent locally.

Both Henderson & Hodge and Clements *et al* remarked on how notoriously secretive this species is during the breeding season and how this resulted in a high proportion of records that referred only to possible breeding, and this was reflected locally in the latest atlas. Clements *et al* note that it is assumed that much such tetrads will in fact hold breeding birds.

### Non-breeding distribution

Figure 2 shows the distribution of all records of Sparrowhawk by tetrad.




Figure 2: Distribution of all Sparrowhawk records at Folkestone and Hythe by tetrad

The Sparrowhawk has been recorded from 30 tetrads (97%) and is only absent from TR13 R which is a tiny stretch of shingle beach.

Taylor *et al* stated that during the period when the Sparrowhawk was absent as a breeding bird in Kent there were still almost annual records of migrants and these increased during their review period due to greater observer coverage. Wernham *et al* (2002) describe an autumn passage from northern Europe, through eastern Britain, to wintering grounds in the Netherlands, Belgium and France, with a return movement in spring.

Suspected migrants have been seen locally on many occasions, including birds seen arriving in off the sea or departing out to sea. Often these records involve single birds but up to six have been noted on several occasions, and on the 12<sup>th</sup> October 1998 a total of 13 flew in/west at Abbotscliffe, with a further seven flying in/west there three days later.

An historical winter count of 10 to 15 present at Sandling Park during a cold spell in January 1956 is also worthy of note.



Sparrowhawk at Saltwood Castle (Brian Harper)



Sparrowhawk at Lympe (Phil Sharp)



Sparrowhawk at Saltwood Castle (Brian Harper)



Sparrowhawk at West Hythe (Brian Harper)

### References

Clements, R., Orchard, M., McCanch, N. & Wood, S. 2015. *Kent Breeding Bird Atlas 2008-13*. Kent Ornithological Society

Harrison, J. M. 1953. *The Birds of Kent*. Witherby, London.

Henderson, A. & Hodge, T. 1998. The Kent Breeding Bird Atlas 1988-94. *Kent Bird Report* 1996: 134-272.

Kent Bird Reports. Kent Ornithological Society

Knight, V. and Tolputt, F., 1871. List of birds observed in Folkestone and its immediate neighbourhood, *Folkestone Natural History Society Annual Report*, 1871

Taylor, D., Davenport, D. & Flegg, J. 1981. *Birds of Kent*. Kent Ornithological Society

Ticehurst, N. F. 1909. *A History of the Birds of Kent*. Witherby, London.

Wernham, C., Toms, M., Marchant, J., Clark, J., Siriwardena, G. & Baillie, S., 2002. *The Migration Atlas: movements of the birds of Britain and Ireland*. British Trust for Ornithology

### **Acknowledgements**

The tetrad map images were produced from the Ordnance Survey [Get-a-map service](#) and are reproduced with kind permission of [Ordnance Survey](#).